


Spanning tight components in 4-uniform hypergraphs



Francesco Di Braccio

Brian Hearn

Joanna Lada

Mihir Neve

Lu-Ming Zhang

London School of Economics, United Kingdom

JCDG³, Tokyo, 10 September 2026

Strong connectivity in hypergraphs

Definition

A k -uniform hypergraph (k -graph) is a hypergraph where all edges contain exactly k vertices.

Strong connectivity in hypergraphs

Definition

A k -uniform hypergraph (k -graph) is a hypergraph where all edges contain exactly k vertices.

Question

What degree conditions guarantee “strong” connectivity in a k -graph?

Strong connectivity in hypergraphs

Definition

A k -uniform hypergraph (k -graph) is a hypergraph where all edges contain exactly k vertices.

Question

What degree conditions guarantee “strong” connectivity in a k -graph?

Example

- Dirac: Minimum degree $n/2 \implies$ Hamiltonicity.

Spanning subgraphs and tight connectivity

Definition

- A subgraph is **spanning** if it touches every vertex.

Spanning subgraphs and tight connectivity

Definition

- A subgraph is **spanning** if it touches every vertex.
- A subgraph is **tightly connected** if there is a **tight walk** between every two edges.




Figure: A tight walk.

Spanning subgraphs and tight connectivity

Definition

- A subgraph is **spanning** if it touches every vertex.
- A subgraph is **tightly connected** if there is a **tight walk** between every two edges.




Figure: A tight walk.

Spanning subgraphs and tight connectivity

Definition

- A subgraph is **spanning** if it touches every vertex.
- A subgraph is **tightly connected** if there is a **tight walk** between every two edges.




Figure: A tight walk.

Spanning subgraphs and tight connectivity

Definition

- A subgraph is **spanning** if it touches every vertex.
- A subgraph is **tightly connected** if there is a **tight walk** between every two edges.




Figure: A tight walk.

Spanning subgraphs and tight connectivity

Definition

- A subgraph is **spanning** if it touches every vertex.
- A subgraph is **tightly connected** if there is a **tight walk** between every two edges.




Figure: A tight walk.

Spanning subgraphs and tight connectivity

Definition

- A subgraph is **spanning** if it touches every vertex.
- A subgraph is **tightly connected** if there is a **tight walk** between every two edges.




Figure: A tight walk.

Spanning subgraphs and tight connectivity

Definition

- A subgraph is **spanning** if it touches every vertex.
- A subgraph is **tightly connected** if there is a **tight walk** between every two edges.




Figure: A tight walk.

Spanning subgraphs and tight connectivity

Definition

- A subgraph is **spanning** if it touches every vertex.
- A subgraph is **tightly connected** if there is a **tight walk** between every two edges.
- A **tight component** is an edge-maximal tightly connected subgraph.




Figure: A tight walk.

Spheres in hypergraphs

“Definition”

A $(k - 1)$ -*sphere* in a k -graph is a tightly connected subgraph “homeomorphic” to a $(k - 1)$ -sphere.

Example: a spanning 1-sphere in a graph is a Hamilton cycle.

Spheres in hypergraphs

“Definition”

A $(k - 1)$ -*sphere* in a k -graph is a tightly connected subgraph “homeomorphic” to a $(k - 1)$ -sphere.

Example: a spanning 1-sphere in a graph is a Hamilton cycle.

Question (Conlon, Gowers)

What degree conditions force a spanning $(k - 1)$ -sphere in a k -graph?

Codegree

Definition

The **codegree** of a $(k - 1)$ -tuple T is the number of edges containing T .




Figure: A triple of codegree 5.

Minimum codegree and spanning spheres

Theorem (Georgakopoulos, Haslegrave, Montgomery, Narayanan (2022))

*Every n -vertex 3-graph with **minimum codegree** at least $n/3 + o(n)$ contains a spanning 2-sphere.*

Conjecture (Georgakopoulos, Haslegrave, Montgomery, Narayanan (2022))

*Every n -vertex k -graph with **minimum codegree** at least n/k contains a spanning $(k - 1)$ -sphere.*

Minimum codegree and spanning spheres

Theorem (Georgakopoulos, Haslegrave, Montgomery, Narayanan (2022))

Every n -vertex 3-graph with *minimum codegree* at least $n/3 + o(n)$ contains a spanning 2-sphere.

Conjecture (Georgakopoulos, Haslegrave, Montgomery, Narayanan (2022))

Every n -vertex k -graph with *minimum codegree* at least n/k contains a spanning $(k - 1)$ -sphere.

Remark

GHMN constructed k -graphs with

- *minimum codegree* $\approx n/k$,
- *no spanning* $(k - 1)$ -spheres,
- *no spanning tight components*.

Minimum codegree and spanning tight components

Conjecture (Illingworth, Lang, Müyesser, Parczyk, Sgueglia (2025))

Every n -vertex k -graph with minimum codegree at least $\lfloor n/k \rfloor$ has a spanning tight component.

Minimum codegree and spanning tight components

Conjecture (Illingworth, Lang, Müyesser, Parczyk, Sgueglia (2025))

*Every n -vertex k -graph with minimum codegree at least $\lfloor n/k \rfloor$ has a **spanning tight component**.*

Remark

- $k = 2$: *trivial*.
- $k = 3$: *a short argument (Cooley, Mycroft).*

Minimum codegree and spanning tight components

Conjecture (Illingworth, Lang, Müyesser, Parczyk, Sgueglia (2025))

*Every n -vertex k -graph with minimum codegree at least $\lfloor n/k \rfloor$ has a **spanning tight component**.*

Remark

- $k = 2$: *trivial*.
- $k = 3$: *a short argument (Cooley, Mycroft).*


Theorem (Di Braccio, H., Lada, Neve, Zhang (2026+))

*Every n -vertex 4-graph with minimum codegree at least $\lfloor n/4 \rfloor$ has a **spanning tight component**.*

$\lfloor n/4 \rfloor$ is tight

Construction (Georgakopoulos,
Haslegrave, Montgomery, Narayanan (2022))

Equipartition $V = V_0 \sqcup V_1 \sqcup V_2 \sqcup V_3$.
For each $v \in V_i$, set $v' := i$.



$\lfloor n/4 \rfloor$ is tight


Construction (Georgakopoulos,
Haslegrave, Montgomery, Narayanan (2022))

Equipartition $V = V_0 \sqcup V_1 \sqcup V_2 \sqcup V_3$.

For each $v \in V_i$, set $v' := i$.

Add edges $abcd$ such that:

- $a' + b' + c' + d' \equiv 1 \pmod{4}$



$\lfloor n/4 \rfloor$ is tight


Construction (Georgakopoulos,
Haslegrave, Montgomery, Narayanan (2022))

Equipartition $V = V_0 \sqcup V_1 \sqcup V_2 \sqcup V_3$.

For each $v \in V_i$, set $v' := i$.

Add edges $abcd$ such that:

- $a' + b' + c' + d' \equiv 1 \pmod{4}$



Remark

Every tight component avoids some V_i .

$\lfloor n/4 \rfloor$ is tight


Construction (Georgakopoulos, Haslegrave, Montgomery, Narayanan (2022)[†])

Equipartition $V = V_0 \sqcup V_1 \sqcup V_2 \sqcup V_3$.

For each $v \in V_i$, set $v' := i$.

Add edges $abcd$ such that either:

- $a' + b' + c' + d' \equiv 1 \pmod{4}$
- † $a' \equiv b' \equiv c' - 1 \equiv d' - 1 \pmod{4}$.



Remark

Every tight component avoids some V_i .

Proof outline

What's simpler than a 4-graph?

Proof outline

What's simpler than a 4-graph?

A 3-graph!

Proof outline

Preprocessing

- Each *triple* (of vertices) is contained in a unique tight component.

Proof outline

Preprocessing

- Each *triple* (of vertices) is contained in a unique tight component.
- Assign a colour to each tight component.
- Colour each triple according to the colour of its tight component.
(Obtain an *edge-colouring* of the complete 3-graph $K_n^{(3)}$.)

Proof outline

Preprocessing

- Each *triple* (of vertices) is contained in a unique tight component.
- Assign a colour to each tight component.
- Colour each triple according to the colour of its tight component.
(Obtain an *edge-colouring* of the complete 3-graph $K_n^{(3)}$.)
- If any vertex misses two colours, then *merge* these colours.

Proof outline

Preprocessing

- Each *triple* (of vertices) is contained in a unique tight component.
- Assign a colour to each tight component.
- Colour each triple according to the colour of its tight component.
(Obtain an *edge-colouring* of the complete 3-graph $K_n^{(3)}$.)
- If any vertex misses two colours, then *merge* these colours.
- **Main point:** *Spanning colour* $\implies$ *Spanning tight component*.

Proof outline

Preprocessing

- Each *triple* (of vertices) is contained in a unique tight component.
- Assign a colour to each tight component.
- Colour each triple according to the colour of its tight component. (Obtain an *edge-colouring* of the complete 3-graph $K_n^{(3)}$.)
- If any vertex misses two colours, then *merge* these colours.
- **Main point:** Spanning colour $\implies$ Spanning tight component.

Outline of the proof

Show that

- there are ≤ 6 colours,
- if there is no spanning tight component, then there are ≥ 7 colours.

There are ≤ 6 colours

What's simpler than a 3-graph?

There are ≤ 6 colours

What's simpler than a 3-graph?

A graph!

There are ≤ 6 colours

Lemma

Any edge-colouring of K_n where every edge lies in $\geq \lfloor (n+1)/4 \rfloor$ monochromatic triangles uses ≤ 5 colours.


There are ≤ 6 colours

Lemma

Any edge-colouring of K_n where every edge lies in $\geq \lfloor (n+1)/4 \rfloor$ monochromatic triangles uses ≤ 5 colours.

Why is this lemma useful?

Fix a vertex u of $K_n^{(3)}$, and consider its *link graph* $L(u)$.



There are ≤ 6 colours

Lemma

Any edge-colouring of K_n where every edge lies in $\geq \lfloor (n+1)/4 \rfloor$ monochromatic triangles uses ≤ 5 colours.

Why is this lemma useful?

*Fix a vertex u of $K_n^{(3)}$, and consider its **link graph** $L(u)$.
There are ≤ 5 colours in $L(u)$.*

There are ≤ 6 colours

Lemma

Any edge-colouring of K_n where every edge lies in $\geq \lfloor (n+1)/4 \rfloor$ monochromatic triangles uses ≤ 5 colours.

Why is this lemma useful?

*Fix a vertex u of $K_n^{(3)}$, and consider its **link graph** $L(u)$.*

There are ≤ 5 colours in $L(u)$.

Since u misses at most 1 colour, there are ≤ 6 colours in $K_n^{(3)}$.

Open problems

Conjecture (GHMN (2022), ILMPS (2025))

Every n -vertex k -graph with minimum codegree at least n/k contains a

- *spanning tight component (ILMPS, 2025),*
- *spanning $(k - 1)$ -sphere (GHMN, 2022).*

Open problems

Conjecture (GHMN (2022), ILMPS (2025))

Every n -vertex k -graph with minimum codegree at least n/k contains a

- *spanning tight component (ILMPS, 2025),*
- *spanning $(k - 1)$ -sphere (GHMN, 2022).*

Question

Can we bound the number of colours in an edge-colouring of K_n where every edge lies in at least $f(n)$ monochromatic triangles?